

**27th INTERNATIONAL CHEMISTRY OLYMPIAD  
PRACTICAL EXAMINATION****FRIDAY, JULY 14, 1995**

*Please read the entire procedure and the Student's Report before beginning the experiment.*

**WARNING:** You must wear your safety goggles in the Laboratory and use the pipette bulbs provided. Should you remove your goggles, or pipette by mouth, you will be given a warning. A second warning will result in a five point penalty. A third warning will result in removal from the laboratory. Removal from the laboratory will result in a zero mark for the practical examination.

**NOTES:**

- 1) Write your name, student number (see student's No. on the table) and the delegation at the top of each student's report sheet.
- 2) Begin only after the instructor has given a **START** message.
- 3) You will be given 5 hours to finish the whole practical exam, including writing your report.
- 4) All the answers are to be written on the student's report sheets within the blanks provided. Only those answers given on the correct positions will be taken into consideration.
- 5) Write your report with the ballpen provided.
- 6) Use deionized water except for cooling.
- 7) The significant figures should be used properly.

*Ar of some elements*

|           |              |           |              |
|-----------|--------------|-----------|--------------|
| <b>H</b>  | <b>1.008</b> | <b>S</b>  | <b>32.06</b> |
| <b>C</b>  | <b>12.01</b> | <b>K</b>  | <b>39.10</b> |
| <b>N</b>  | <b>14.01</b> | <b>Cu</b> | <b>63.54</b> |
| <b>O</b>  | <b>16.00</b> | <b>I</b>  | <b>126</b>   |
| <b>Na</b> | <b>22.99</b> |           |              |

**PRACTICAL PROBLEM I****Identification of Unknown Solutions****Reagents**

|  |  |
|--|--|
| <b>H<sub>2</sub>SO<sub>4</sub> (conc.)</b>       | <b>H<sub>2</sub>SO<sub>4</sub> (6 mol·dm<sup>-3</sup>)</b>     |
| <b>HNO<sub>3</sub>(conc.)</b>                    | <b>HNO<sub>3</sub> (6 mol·dm<sup>-3</sup>)</b>                 |
| <b>HCl(conc.)</b>                                | <b>HCl (6 mol·dm<sup>-3</sup>)</b>                             |
| <b>Ba(OH)<sub>2</sub>(satd.)</b>                 | <b>NaOH (6 mol·dm<sup>-3</sup>)</b>                            |
| <b>BaCl<sub>2</sub>(0.5 mol·dm<sup>-3</sup>)</b> | <b>Ba(NO<sub>3</sub>)<sub>2</sub>(0.5 mol·dm<sup>-3</sup>)</b> |

**Desk equipments**

One test tube holder  
five small test tubes

**Problem**

You are supplied with five different solutions contained in five test tubes labeled as A,B,C,D and E, respectively. The solution, in each test tube, contains one of the following compounds



**Identify these solutions.**

**NOTES:**

- (1) You can only select the provided reagents and use a procedure as simple as possible to complete your task. You are getting a mark not only according to the correct identification, but also to the number of steps you have taken.
- (2) You have to carry out the whole analysis by using the provided amount of these unknown solutions. Supplement of them will be available, but it will reduce the mark you obtain.

## PRACTICAL PROBLEM II

### Preparation of cis-Copper-bis-Glycinate Hydrate [Cu(gly)<sub>2</sub>·xH<sub>2</sub>O]

Copper(II) amino acidate coordination compounds are monomeric units for synthesizing important biopolymers such as metalloenzymes like ceruloplasmin, on which every living organism depends. In laboratory cis-copper-bisglycinate hydrate can be produced by the reaction of cupric hydroxide with glycine at a temperature of *ca.* 70°C.

#### Reagents:

CuSO<sub>4</sub>·5H<sub>2</sub>O(s)

NH<sub>3</sub>·H<sub>2</sub>O( 3 mol·dm<sup>-3</sup>)

glycine(s)

95% ethanol,

acetone

NaOH( 2 mol·dm<sup>-3</sup>)

BaCl<sub>2</sub>( 0.5 mol·dm<sup>-3</sup>)

#### Desk equipments

beaker 250 cm<sup>3</sup> ×4

graduated cylinder 100cm<sup>3</sup> ×1

filter flask (shared)

Buchner funnel 60mm ×1

watch glass ×2

dropper ×2

spatula ×1

glass stirring rod ×1

aspirator

thermometer ( at least 100° C)

#### Procedure:

##### 1. Preparation of Cu(OH)<sub>2</sub>

(1) Dissolve your pre-weighted sample of CuSO<sub>4</sub>·5H<sub>2</sub>O (6.0 g) in 40 cm<sup>3</sup> of water with a 250 cm<sup>3</sup> beaker as a container.

(2) Add slowly 3 mol·dm<sup>-3</sup> ammonia solution to the CuSO<sub>4</sub> solution, gently stirring, until the precipitate is completely dissolved and the solution is turning blue-violet.

(3) Add 2 mol·dm<sup>-3</sup> NaOH solution to the above solution until no more precipitate formed.

(4) Filter the precipitate over a Buchner funnel under reduced pressure. Wash the precipitate with water until no  $\text{SO}_4^{2-}$  ion is detected in the filtrate.

(5) Collect  $\text{Cu}(\text{OH})_2$  for the preparation of  $\text{Cu}(\text{gly})_2 \cdot x\text{H}_2\text{O}$ .

**Write the equations for the main chemical reactions having taken place in the above procedure.**

## 2. Preparation of cis-copper-bisglycinate hydrate

(1) Dissolve a pre-weighted sample of glycine (3.6 g) in  $130 \text{ cm}^3$  of water and then warm the solution in a hot water bath ( $70^\circ\text{C}$ ). Add the  $\text{Cu}(\text{OH})_2$  to the solution, gently stirring until the precipitate is dissolved. Perform a hot filtration and add  $10 \text{ cm}^3$  of 95% ethanol.

(2) Cool the solution and then needle-like crystals appear, place it in the ice water bath for 10 min.

(3) Filter the crystals over a Buchner funnel under reduced pressure, wash once with  $10 \text{ cm}^3$  of ethanol-water mixing solvent and then twice with  $10 \text{ cm}^3$  acetone, squeeze the crystals as dry as possible on the funnel.

(4) Collect the crystals to a watch glass and dry it (consult your supervisor).

(5) Half an hour later weigh the product. Write the mass of product and the percentage of yield on your student's report. Give the expressions for calculation to show how you calculate.

## PRACTICAL PROBLEM III

### Determination of copper(II) content in $\text{Cu}(\text{gly})_2 \cdot x\text{H}_2\text{O}$

The Cu(II) content in  $\text{Cu}(\text{gly})_2 \cdot x\text{H}_2\text{O}$  crystals prepared yourself can be determined by iodometry with starch solution as indicator. Based on the data obtained one can calculate the moles of hydrates of crystals in  $\text{Cu}(\text{gly})_2 \cdot x\text{H}_2\text{O}$ .

#### Reagents

Standard  $\text{KIO}_3$  (see the label on the bottle to get the accurate concentration)

$\text{H}_2\text{SO}_4$  (1.0 mol · dm<sup>-3</sup>) as indicator.

KI (0.6 mol · dm<sup>-3</sup>)

KSCN (2 mol · dm<sup>-3</sup>)

Starch (0.5%)

$\text{Na}_2\text{S}_2\text{O}_3$  (to be standardized)

#### Desk equipments

buret 50cm<sup>3</sup> ×1

pipette 25cm<sup>3</sup> ×1

pipette bulb

beakers (dry) 100cm<sup>3</sup> ×2

volumetric flask 100cm<sup>3</sup> ×1

Erlenmeyer flask 250 cm<sup>3</sup> ×3

graduated cylinder 10cm<sup>3</sup> ×3, 100cm<sup>3</sup> ×1

wash bottle

single pan balance (shared)

hot water bath (shared)

#### Procedure

1. Standardization of  $\text{Na}_2\text{S}_2\text{O}_3$  solution

(1) Transfer 25.00 cm<sup>3</sup> of standard  $\text{KIO}_3$  solution to an Erlenmeyer flask.

(2) Add 5 cm<sup>3</sup> of water, 10 cm<sup>3</sup> of KI solution and 5 cm<sup>3</sup> of  $\text{H}_2\text{SO}_4$  (1.0 mol · dm<sup>-3</sup>) to the flask.

(3) Titrate immediately with  $\text{Na}_2\text{S}_2\text{O}_3$  solution.

(4) Add 2 cm<sup>3</sup> starch solution when the colour of the titrand turns pale yellow.

(5) Continue titrating until the blue color of the solution disappears.

(6) Proceed with step (1)—(5) twice parallelly.

**2. Determination of Cu(II) content in  $\text{Cu}(\text{gly})_2 \cdot x\text{H}_2\text{O}$** 

- (1) Weigh 1.0—1.2 g (precision of  $\pm 0.0002$  g) of  $\text{Cu}(\text{gly})_2 \cdot x\text{H}_2\text{O}$  with a dry  $100 \text{ cm}^3$  beaker as the container.**
- (2) Dissolve it with  $40 \text{ cm}^3$  of water and  $8 \text{ cm}^3$  of  $\text{H}_2\text{SO}_4$  ( $1.0 \text{ mol} \cdot \text{dm}^{-3}$ ).**
- (3) Transfer the above solution quantitatively to a  $100 \text{ cm}^3$  volumetric flask and dilute to the mark.**
- (4) Transfer  $25.00 \text{ cm}^3$  of the Cu(II) Solution to an Erlenmeyer flask, add  $50 \text{ cm}^3$  of water and  $10 \text{ cm}^3$  of KI solution to the flask.**
- (5) Titrate immediately with standardized  $\text{Na}_2\text{S}_2\text{O}_3$  solution.**
- (6) Add  $2 \text{ cm}^3$  of starch solution and  $3 \text{ cm}^3$  of KSCN solution to the flask when the colour of the titrand turns from brown to pale yellow.**
- (7) Titrate continuously until the blue color of the solution disappears.**
- (8) Proceed with steps (4)—(7) twice parallelly.**